BLS First Responder Unit

| Company: | | Fleet ID: | | | | | |
|------------|----------------|---|-------|------|-----------|--|--|
| VIN: | VIN:Inspector: | | | | | | |
| Inspection | | Needs Decal: Yes/No (Quantity:) | | | | | |
| | | Deficiencies | | | Corrected | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| Line # | Quantity | Description | Pass | Fail | Notes | | |
| Line π | Quantity | General Supplies | 1 455 | 1 an | Notes | | |
| 1 | 1 ea. | Blanket | | 1 | | | |
| 2 | 1 ea. | Obstetrical (OB) Kit (commercially packaged) | | | | | |
| 3 | 2 ea. | Clean linen sheets or Mylar blanket (for burns) | | | | | |
| 4 | 1 ea. | Maryland Triage Kit: Enough triage tags and | | | | | |
| | | color-coded ribbons to triage 25 patients. Kits | | | | | |
| | | must contain paperwork for the treatment and | | | | | |
| | | transportation areas. | | | | | |
| 5 | _ | ANSI 207-2006 class II reflective safety vests | | | | | |
| 6 | 1 00 | for each crew member PHMSA Emergency Response Guidebook | | | | | |
| O | 1 ea. | (ERG): Current edition in either print or | | | | | |
| | | electronic format | | | | | |
| 7 | 1 ea. | Environmental carbon monoxide alarming | | | | | |
| | | device (OPTIONAL) | | | | | |
| | | Portable First Aid Kit | | | | | |
| 8 | 12 ea. | Sterile gauze pads (min. 4" x 4") | | | | | |
| 9 | 4 ea. | Sterile dressings (min. 5" x 9") | | | | | |
| 10 | 1 ea. | Hemostatic impregnated dressing | | | | | |
| | | (OPTIONAL): Impregnated with either | | | | | |
| | | chitosan or kaolin. Form of either roller gauze or trauma dressings (2" x 2" and/or 4" x 4" | | | | | |
| | | dressings are not acceptable. | | | | | |
| 11 | 4 rolls | Self-adhering gauze bandages (various sizes 2" - 6") | | | | | |
| 12 | 4 ea. | Cravats (triangular bandages): Minimum size is | | | + | | |
| 12 | 4 ca. | 36" x 36". | | | | | |
| 13 | 2 ea. | Cold packs | | | | | |
| 14 | 1 ea. | Bottle normal saline and/or sterile water | | | | | |
| | | (500cc): (check exp. date) | | | | | |
| 15 | 4 ea. | Commercially available tourniquet capable of | | | | | |
| | | stopping arterial blood flow | 1 | | | | |
| 16 | 2 rolls | 1" medical tape (hypoallergenic tape must be | | | | | |
| | I | available) | | 1 | | | |

17

18

19

20

1 ea.

1 ea.

1 ea.

1 ea.

Ring cutter

Bandage scissors or rescue shears (at least 5.5")

Penlight (narrow beam flashlight acceptable)

Stethoscope (must be pediatric-capable)

| | | BLS First Responder Unit |
|------|------------------|---|
| | | Portable First Aid Kit (continued) |
| 21 | 1 ea. | Glucometer kit – must include lancets, test |
| | | strips, alcohol wipes, and band-aids. (check |
| - 22 | | exp. date on test strips.) |
| 22 | 1 ea. | Adult Regular BP Cuff |
| 23 | 1 ea. | Adult Large BP Cuff Child BP Cuff |
| 25 | 1 ea. 6 pairs | Non-latex exam gloves (assorted sizes) |
| 26 | 1 ea. | Portable sharps container |
| 27 | 1 ea. | Pulse oximeter (OPTIONAL)(Adult and Peds |
| 27 | ı cu. | sensors recommended) |
| 28 | 1 ea. | Clean kit large enough to carry all supplies in |
| | | "Portable First Aid Kit" |
| | | Medications and Delivery Devices (check expiration dates) |
| 29 | 5mg | Albuterol sulfate- may substitute one albuterol |
| | | hand-held inhaler |
| 30 | 325mg | Aspirin (chewable) |
| 31 | 2 ea. | Glucose paste tubes |
| 32 | 6ea. | Isopropyl Alcohol Wipes |
| 33 | 4mg | Naloxone (Narcan) |
| 34 | 2 ea. | Intranasal medication delivery device (not |
| | 2 0 | necessary if pre-packaged with naloxone) |
| 35 | 1 ea. | Epinephrine auto-injector (adult): Draw-up epi |
| | | may be substituted if jurisdiction is approved |
| | | for the optional protocol |
| 36 | 1 ea. | Epinephrine auto-injector (pediatric): Draw-up |
| | | epi may be substituted if jurisdiction is |
| | | approved for the optional protocol |
| 37 | - | DuoDote (OPTIONAL) |
| | | AED |
| 38 | 1 ea. | AED – pediatric cable |
| 39 | 2 ea. | AED pads (adult) – (check exp. dates) |
| 40 | 1 ea. | AED pads (pediatric) – (check exp. date)(Only |
| 10 | ı cu. | required if device or pads does not have dose |
| | | attenuator) |
| 41 | 1 ea. | Spare battery (if required) |
| 42 | 1 ea. | Razor |
| 43 | 1 ea. | Washcloth or towel (OPTIONAL) |
| 7-3 | ı ca. | Biohazard Items |
| 44 | 1ea. | Surgical masks: Should be provided for each |
| | ıca. | seated position on the unit. Must have one for |
| | | patient. |
| 45 | 1ea. | Gowns (impenetrable to body fluids): Should |
| 45 | ıca. | be provided for each seated position on the |
| | | unit. |
| 4.0 | 1 | |
| 46 | 1ea. | Eye/facial shield (may be combined with |
| | | mask): Should be provided for each seated position on the unit. |
| | | |
| 47 | 1ea. | Particulate respirator N95 or greater for each |
| | | crew member N95 needs to be fit tested for a |
| | | proper fit. Should be provided for each seated |
| | | position on the unit. |

| | | BLS First Responder Unit | |
|----|----------------|---|--|
| | | Biohazard Items (continued | |
| 48 | 1 ea. | Appropriate disinfectant: Effective against | |
| | | pathogens to include hepatitis B virus (HBV), | |
| | | human immunodeficiency virus (HIV), and M. | |
| | | tuberculosis (TB), and SARS-CoV2. Minimum | |
| | | of wipes are sufficient. (check exp. date) | |
| 49 | 2 ea. | Nasal cannula (adult) | |
| 50 | 2 ea. | Nasal cannula (pediatric) | |
| 51 | 2 ea. | Non-rebreather mask (adult) | |
| 52 | 2 ea. | Non-rebreather mask (pediatric) | |
| 53 | 1 ea. | Nebulizers - not necessary if using hand-held | |
| | | albuterol inhaler | |
| 54 | 1 ea. | Adult (1,000 – 1,200 mL) hand-operated, self | |
| | | re-expanding bag resuscitator without a pop-off | |
| | | valve or with a selectable pop-off valve. Peep | |
| | | valve (REQUIRED) and viral filter recommended. | |
| | | | |
| | | An oxygen inlet Reservoir tube | |
| 55 | 1 00 | Transparent face mask (adult) | |
| 56 | 1 ea. 1 ea. | Child (750 mL) hand-operated, self re- | |
| 30 | i ea. | expanding, bag resuscitator without a pop-off | |
| | | valve or with a selectable pop-off valve. PEEP | |
| | | valve (REQUIRED) and viral filter | |
| | | recommended. | |
| | | An oxygen inlet | |
| | | Reservoir tube | |
| 57 | 1 ea. | Infant (450 - 500 mL) hand-operated, self re- | |
| | ı cu. | expanding, bag resuscitator without a pop-off | |
| | | valve or with a selectable pop-off valve. PEEP | |
| | | valve (REQUIRED) and viral filter | |
| | | recommended. | |
| | | An oxygen inlet | |
| | | Reservoir tube | |
| 58 | 1 set | Transparent face mask (neonate through small | |
| | | adult; a set is 4 sizes): Masks can be included | |
| | | with BVM | |
| 59 | 1 set | Oropharyngeal airway (newborn through large | |
| | | adult; a set is 6 sizes) | |
| 60 | 1 set | Nasopharyngeal airways (18 Fr – 34 Fr., a set | |
| | | is 6 sizes): (check exp. dates) | |
| 61 | 1 ea. | Water soluble lubricant (1 tube or 3 packets): | |
| | | (check exp. dates) | |
| 62 | 1 ea. | Medical oxygen cylinder with at least 300 L | |
| | | capacity (required "E," "D," or super D size): | |
| | | Portable tank minimum of 500 psi. Portable | |
| | | tank must be in a portable bag that can be | |
| | | carried to patient side. | |
| | | | |
| | | (#1) Year PSI | |
| | | Cylinder properly color-coded (green = steel, | |
| | | unpainted = brushed metal for aluminum or | |
| | | stainless steel) | |
| | | Free of grease, oil, or other flammable organic | |
| | | material | |
| L | <u> </u> | 1 | |

| | | BLS First Responder Unit |
|-----|-------|--|
| | | Portable Oxygen Kit (continued) |
| | | Passed hydrostatic testing within the past 5 years for aluminum and steel cylinders: Steel cylinders with a hydrostatic test date followed |
| | | by a star is acceptable for 10 years. Regulator shall have a pressure gauge to indicate the pressure of oxygen remaining in |
| | | the cylinder (not gravity-dependent). Can be separate or combined with oxygen pressure gauge. |
| | | A variable flow valve and a flowmeter capable of delivering at least 15 LPM, with a dial-down rate to a minimum of 2 LPM |
| | | Accurate within 1 LPM when setting equal to or less than 5 LPM Test reading ofLPM when flowmeter set at 4 LPM (3 – 5 LPM) |
| | | Accurate within 1.5 LPM when setting between 6 and 10 LPM Test reading ofLPM when flowmeter set at 10 LPM (8.5 – 11.5 LPM) |
| | | Accurate within 2 LPM when setting equal to or greater than 15 LPM Test reading of _LPM when flowmeter set at 15 LPM (13 – 17 LPM) |
| 63 | 1 ea. | Clean kit large enough to carry all supplies in "Portable Oxygen Kit" |
| | | Portable Suction Unit |
| 64 | 1 ea. | Manual pumps must meet the same testing requirements as a battery-operated suction device, and have the following: |
| 64a | | All of the required manufacturer parts |
| 64b | | Adult soft tip catheter |
| 64c | | Adult hard tip catheter |
| 64d | | Pediatric soft catheter |
| 65 | | If using battery-powered suction unit, it must be capable of operating continuously under suction for at least 20 minutes with a rigid suction tip |
| 65a | | Must be able to develop 11.81 inches of water vacuum (300 mmHg) within 4 seconds of clamping Test reading at 4 secin/Hg |
| 65b | | A free air flow of at least 20 LPM at the end of the suction tube. Test readingLPM |
| 65c | - | Assorted catheters 6 – 18 Fr and rigid suction tips: One must be between 6 and 12 Fr AND one between 12 and 18 Fr (check exp. dates) |
| | | First Responder Unit Vehicle |
| 66 | | Functional emergency warning lights Functional emergency audible warning devices (not horn) |
| 68 | | Functional head, tail, and signal lights |

| BLS First Responder Unit | | | | | |
|--|-------|--|--|--|--|
| First Responder Unit Vehicle (continued) | | | | | |
| 69 | 1 ea. | Fire extinguisher (5 lb. multipurpose dry | | | |
| | | chemical). Check label or bottom of cylinder | | | |
| | | for date. Must be mounted or secured. May be | | | |
| | | mounted in outside compartment. | | | |
| 70 | 3 ea. | Reflective road hazard triangles or 3 small | | | |
| | | traffic cones (Flares are not an acceptable | | | |
| | | substitute) | | | |
| 71 | 1 ea. | Radio w/ capability to communicate with | | | |
| | | PSAP/EMRC | | | |